

The National Immunisation Schedule

Age	Disease to protect against	Vaccine
Pregnancy	Tetanus + diphtheria + whooping cough (pertussis)	Boostrix® (28 to 38 weeks)
	Influenza	Any time during pregnancy.
6 Weeks	Rotavirus (first dose must be given before 15 weeks)	Rotarix® (oral)
	Diphtheria + tetanus + whooping cough (pertussis) + polio + hepatitis B + <i>Haemophilus influenzae</i> type b (Hib)	Infanrix® hexa
	Pneumococcal disease	Synflorix®
3 Months	Rotavirus (second dose must be given before 25 weeks)	Rotarix® (oral)
	Diphtheria + tetanus + whooping cough + polio + hepatitis B + <i>Haemophilus influenzae</i> type b (Hib)	Infanrix® hexa
	Pneumococcal disease	Synflorix®
5 Months	Diphtheria + tetanus + whooping cough + polio + hepatitis B + <i>Haemophilus influenzae</i> type b (Hib)	Infanrix® hexa
	Pneumococcal disease	Synflorix®
15 Months	<i>Haemophilus influenzae</i> type b (Hib)	Hiberix®
	Measles + mumps + rubella	Priorix®
	Pneumococcal disease	Synflorix®
	Varicella (chickenpox)	Varilrix®
4 Years	Diphtheria + tetanus + whooping cough + polio	Infanrix® IPV
	Measles + mumps + rubella	Priorix®
11+12 Years	Tetanus + diphtheria + whooping cough	Boostrix®
	Human papillomavirus (HPV)	Gardasil®9 (2 doses, 6 months apart)
45 Years	Tetanus + diphtheria	ADT® Booster
65 Years	Tetanus + diphtheria	ADT® Booster
	Influenza	Given annually.